

Original Article

# Do We Need Fluid Restriction after Stapled Hemorrhoidopexy? A Pilot, Double-blinded, Randomized Controlled Trial

Chun-Wei Yu<sup>1</sup>  
Nien-Ying Tsai<sup>2</sup>  
Shu-Wen Jao<sup>1</sup>  
Chao-Yang Chen<sup>1</sup>  
Chia-Cheng Wen<sup>1</sup>  
Chien-Chang Kao<sup>3</sup>  
Kuan-Ling Lin<sup>1</sup>  
Po-Hsien Wu<sup>1</sup>  
Yi-Chiao Cheng<sup>1</sup>

<sup>1</sup>Division of Colon and Rectal Surgery,  
Department of Surgery,

<sup>2</sup>Department of General Medicine,

<sup>3</sup>Division of Urological Surgery, Department  
of Surgery, Tri-Service General Hospital,  
National Defense Medical Center, Taipei,  
Taiwan

## Key Words

Stapled hemorrhoidopexy;  
Fluid restriction;  
Urinary retention

**Background.** Conventional hemorrhoidectomy leads to a high urinary retention rate and fluid restriction is commonly recommended to minimize complications. However, the need for postoperative fluid restriction among patients who have undergone stapled hemorrhoidopexy is unclear. We aimed to determine whether fluid restriction after stapled hemorrhoidopexy with/without partial external hemorrhoidectomy could reduce urinary retention and postoperative pain.

**Patients and Methods.** In this prospective, double-blinded, randomized controlled trial, we enrolled patients who had grade III or IV hemorrhoids and underwent stapled hemorrhoidopexy with/without partial external hemorrhoidectomy; 250 mL/12 h of sterile 0.9% saline was administered to the fluid restriction group after the operation, and 1000 mL/12 h was given to the non-fluid-restriction group. We focused on the need for fluid restriction after stapled hemorrhoidopexy.

**Results.** Fifty patients were divided into two groups. The surgical outcomes were of no difference between the two groups. The median subjective pain scores at 0, 8, 16, 24 h after the operation showed no significant difference between the two groups ( $p = 0.55$  at 0 h;  $p = 0.38$  at 8 h;  $p = 0.98$  at 16 h; and  $p = 0.66$  at 24 h). The mean times to first urination after the operation were  $700.04 \pm 455.03$  mins in the non-fluid-restriction group and  $737.16 \pm 426.32$  mins in the fluid-restriction group (no significant difference;  $p = 0.67$ ). No postoperative urinary retention was found in either group of patients.

**Conclusions.** Fluid restriction after stapled hemorrhoidopexy is unnecessary. [*J Soc Colon Rectal Surgeon (Taiwan) 2022;33:41-49*]

**H**emorrhoidal disease, a common anorectal disorder, is the main cause of rectal bleeding, discomfort, and pain for affected patients. The prevalence of hemorrhoid disease has been estimated at 4.4-12.8% in normal adult populations and about 40% in patients with symptoms of anal disease. However, it is difficult to estimate the exact incidence of hemorrhoid disease,

because of patient embarrassment, fear, or discomfort related to the treatment.<sup>1,2</sup>

There are several possible surgical treatments. Conventional hemorrhoidectomy (CH), which is currently considered the most effective treatment, involves excision of the hemorrhoidal cushions. However, it is often accompanied by postoperative pain

Received: January 2, 2022.

Accepted: March 1, 2022.

Correspondence to: Dr. Yi-Chiao Cheng, Division of Colon and Rectal Surgery, Department of Surgery, Tri-Service General Hospital, National Defense Medical Center, No. 325, Section 2, Cheng-Kong Rd., Neihu District, Taipei 114, Taiwan. Tel: 886-2-8792-7209; Fax: 886-2-8792-7411; E-mail: ndmcjoe@gmail.com

and disability, which makes it an unpopular choice with patients. The most common complication after CH is acute urinary retention, with an incidence of about 21.9%.<sup>3</sup> Urinary retention causes discomfort, and urethral injury, and urinary tract infection might occur as a result of catheterization. These may also cause delayed discharge and possibly increased medical expenses to the patient.<sup>4,5</sup> Many different approaches have been tried to prevent urinary retention, such as pain control, warm water sitz bath, or restriction of perioperative fluid intake.<sup>6</sup> In 1976, a report revealed that fluid restriction could reduce the incidence of urinary retention following operations for benign anorectal diseases. According to that research, only 3.5% of patients with fluid restriction presented with postoperative urinary retention, but 14.9% of patients without fluid restriction showed it.<sup>4,6</sup>

Stapled hemorrhoidopexy (SH) was first described by Longo in 1998.<sup>7</sup> In contrast to the traditional approach, which involves removing the hemorrhoidal tissue, SH removes the submucosa above the dentate line with a circular stapler. With this position of excision, postoperative pain can be reduced, and the sensitivity of the anal canal can be maintained by preserving the transitional epithelium.<sup>8,9</sup> However, the general incidence of complications after SH may be identical to conventional excisional surgery.<sup>10</sup> Despite the low risk, SH might also contribute to urinary retention.<sup>6</sup> A 1.5% urinary retention rate in the first week after being treated with SH was reported in a review of 1107 patients from 12 Italian coloproctological centers.<sup>9</sup>

To avoid urinary retention, we routinely arrange fluid restriction for patients undergoing SH in clinical practice. However, whether restriction of postoperative fluid intake in such cases is necessary is unknown. Therefore, we designed a prospective, randomized, double-blinded study, and aimed to investigate the influence of perioperative fluid restriction with urinary retention after SH.

## Patients and Methods

### Patients

This prospective, double-blinded, randomized con-

trolled trial included patients with grade III or IV hemorrhoids who underwent SH with/without partial external hemorrhoidectomy at the Colorectal Division of Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan, from February 2018 to December 2019.

Exclusion criteria were as follows: patients younger than 20 years or older than 80 years; allergy to nonsteroidal anti-inflammatory drugs; uremia or impaired renal function; pregnant or breast feeding.

Our research nurse assigned the patients to the fluid restriction (experimental group, EG) or non-fluid-restriction groups (control group, CG) at a 1:1 ratio using a computer-generated list of random numbers (Microsoft Excel). Once the patients had been assigned into either group, only the bedside nurses were informed about the allocation.

This study followed the guidelines of the 7th revision of Helsinki Declaration. All participants gave informed consent at hospital admission. This study was reviewed and approved by the Tri-Service General Hospital Institutional Review Board for human subjects (No. 2-106-05-063). This study also had trial registration with clinicalTrials.gov (Identifier: NCT 04459039). The first registration was July 7<sup>th</sup> 2020.

### Primary endpoint

In order to reveal if fluid restriction could avoid urine retention, the study defined the primary endpoint as the first urination after stapled hemorrhoidopexy with/without partial external hemorrhoidectomy.

### Preparation

All patients were admitted the day before the operation. After basic blood tests, they received uroflowmetry and bladder scans for evaluating residual urine to reduce the possible impact of selection bias related to postoperative urinary retention between the two study groups. Since most of the patients recruited in our study were middle age, the patients were given a clear liquid diet and received sodium phosphate solution or polyethylene glycol electrolyte lavage solu-

tion orally on the night before surgery for colonoscopy in order to rule out possible colon lesions.

## Surgery

All the patients were under intravenous general anesthesia using propofol, and administered local anesthesia with mixed agents (15 mL 2% lidocaine, 10 mL 0.5% bupivacaine and 0.3 mL 1:200,000 epinephrine) into the intersphincteric groove. After receiving colonoscopy routinely, all patients underwent SH with/without partial hemorrhoidectomy for cosmetic concern applied by four experienced colorectal surgeons (SWJ, CYC, CCW, and YCC) after being placed in a prone jackknife position. The devices used for SH were PPH® staplers (Ethicon Inc., Cincinnati, OH, USA) or EEA® staplers (Medtronic plc, Fridley, MN, USA) according to the surgeons' preferences. The extent of internal hemorrhoids, numbers of excised external hemorrhoids, distance between the staple and dentate lines, and operation times were recorded.

After the surgical procedure, ketorolac (30 mg) was prescribed routinely immediately after the operation and every 6 h for 1 day. A rescue tramadol acetaminophen tablet (37.5 mg) was given every 6 h if needed. The patients were asked to report a 0-10 visual analogue scale (VAS) pain score at postoperative 0, 8, 16, and 24 h. No warm water sitz (hip) baths were permitted, nor any medicines other than painkillers.

All patients were allowed to consume a soft diet at 6 h after the operation, but their oral fluid intake was restricted to < 400 mL until their first urination. Within 12 h of surgery, the EG received 1000 mL 0.9% sterile saline intravenously (IV), while the CG received 250 mL similarly. The rate of fluid supply was controlled using an infusion pump. The monitor and the saline bag were masked for double-blinding. Bedside nurses recorded the time to first urination in minutes after the operation, and the time of urethral catheterization if acute urinary retention persisted. Acute urinary retention was defined as being when patients were unable to empty their bladder voluntarily and when physical examinations by doctors demonstrated bladder distention.

## Statistical analysis

Categorical data are reported as absolute numbers and percentages while continuous variables are expressed as the mean  $\pm$  standard deviation or the median and interquartile range. Quantitative data were analyzed using the Mann-Whitney nonparametric *U* test. Categorical data were compared using Chi-squared or Fisher's exact tests, as appropriate. Box-and-whisker plots were used to compare the VAS differences at 0, 8, 16, and 24 h after surgery between the EG and CG groups. Scatter plots of the time to first urination after surgery were used to demonstrate the influence of fluid restriction. A *p* value < 0.05 was considered statistically significant. All analyses were performed using IBM SPSS Statistics for Windows (Version 25.0; IBM Corp., Armonk, NY, USA).

## Results


Fifty-three patients underwent SH. Two were excluded before the operation; one was excluded for severe anemia, which would lead to blood transfusion and increase intravenous fluid volume and make an effect on the result in this study; the other patient had huge prolapsed hemorrhoids and the surgeon decided to perform a conventional hemorrhoidectomy. The remaining 51 patients were randomized into two groups. One patient who refused intravenous fluid after the operation was excluded. Therefore, 50 cases were included finally (Fig. 1).

Table 1 reveals the basic characteristics of both groups. The mean age of the patients was  $54.24 \pm 11.72$  years in the non-fluid-restriction group and  $46.40 \pm 13.92$  years in the fluid-restriction group (*p* = 0.028). However, the gender distributions, body mass index, grades of hemorrhoids, uroflowmetry data, and American Society of Anesthesiology class were not significantly different between the groups.

Surgical outcomes are shown in Table 2. The two groups had no difference in terms of the use of partial external hemorrhoidectomy, operation time, the level of the stapling line, intraoperative IV fluid needs, medicine used, time to first urination, the incidence of



CONSORT 2010 Flow Diagram



**Fig. 1.** Before surgery, two of fifty-three were excluded because of severe anemia and massive volume of hemorrhoidal tissue. One of fifty-three was excluded because the patient refused intravenous fluid. Therefore, remained fifty cases were included.

**Table 1.** Characteristics

	Non-fluid restriction (n = 25)	Fluid restriction (n = 25)	<i>p</i> value
Age (years)	54.24 ± 11.72	46.40 ± 13.92	0.028
Male (%)	16 (64)	13 (52)	0.390
BMI (kg/m <sup>2</sup> )	23.06 ± 3.02	23.19 ± 2.96	0.969
Hemorrhoids, grade			
III	22	19	0.289
IV	3	6	
Pre-operative uroflowmetry			
Peak flow	16.60 ± 8.05	19.29 ± 9.25	0.284
Mean flow	9.16 ± 4.72	11.13 ± 5.96	0.233
Residual urine	23.72 ± 48.70	39.36 ± 50.97	0.092
ASA*			0.478
I	4	8	
II	19	16	
III	2	1	

\* ASA class = American Society of Anesthesiologists Classification.

**Table 2.** Outcomes

	Non-fluid restriction (n = 25)	Fluid restriction (n = 25)	p value
With partial external hemorrhoidectomy	12 (19 lumps)	10 (13 lumps)	0.569
Operation time (min)	32.08 ± 8.47	32.88 ± 10.85	0.954
Staple line (mm above the dentate line)	19.0 ± 4.8	19.8 ± 8.2	1.000
Intraoperative IV fluid (mL)	218 ± 87.66	212 ± 89.30	0.712
Use of rescued medicine	3	5	0.702
Time to 1 <sup>st</sup> urination (min)	700.04 ± 455.03	737.16 ± 426.32	0.669
Urinary retention	0	0	N/A
Complications	2 (PPH, delayed bleeding)	0	0.490
Post-operative LOS*	2.64 ± 0.49	2.60 ± 0.65	0.634

\* LOS = length of stay.

urinary retention, complications, or postoperative length of stay.

The median VAS scores of two groups at 0, 8, 16, and 24 h after the operation are shown as box-and-whisker plots in Fig. 2. The median VAS scores did not reach significant difference at any recorded hours ( $p = 0.548$  at 0 h;  $p = 0.382$  at 8 h;  $p = 0.984$  at 16 h; and  $p = 0.657$  at 24 h).


The scatter plot in Fig. 3 reveals individual patients' time to their first urination after the operation. The mean values were  $700.04 \pm 455.03$  min in the non-fluid-restriction group and  $737.16 \pm 426.32$  min in the fluid-restriction group. No significant difference was found between the groups ( $p = 0.669$ ).

## Discussion


Previous studies had demonstrated the contribut-

ing risk factors of postoperative urinary retention, including age, pre-existing neurologic abnormalities, bladder volume on entry to the post-anesthesia care unit, surgical procedure, length of surgery, intraoperative aggressive fluid administration, postoperative pain, and need for postoperative analgesia, or postoperative opioid use.<sup>5,11</sup> The mechanism of postsurgical urinary retention is still not clear. There are some probable causes, including anxiety, anal distention, bladder distention caused by fluid hydration during surgery, irritation or blockade of pelvic nerves, and reflex inhibition of the urinary bladder detrusor muscle arising from pain.<sup>12,13</sup>

Numerous steps have been reported to prevent postsurgical urinary retention, including the use of parasympathomimetic agents, alpha-adrenergic blockers and anxiolytic agents, restriction of perioperative fluid intake, avoidance of anal packing, use of sitz baths, local anesthesia, short-acting anesthesia, and



**Fig. 2.** The median VAS scores of two groups at 0, 8, 16, and 24 h after the operation.



**Fig. 3.** There was no statistically significant difference in the average time to first urination between the two groups.

outpatient surgery.<sup>14-16</sup>

Increased fluid intake might cause overdistension of the bladder and dysfunction of the detrusor muscle. Some studies showed that fluid restriction can prevent postoperative urinary retention effectively.<sup>6,16,17</sup> In 2006, Toyonaga et al. reported that a significant increase in postoperative urinary retention was noted in patients with IV fluids in excess of 1000 mL, and perioperative fluid restriction was recommended for prevention.<sup>15</sup> However, fluid restriction with preoperative patient preparation, including nothing by mouth and the use of laxatives, can cause thirst and discomfort in patients. That is why we aimed to determine whether fluid restriction after SH is necessary.

Unlike previous research, our study demonstrated that patients in the non-fluid-restriction group did not have higher urinary retention rates than the fluid-restriction group, although 22 of the 50 patients had external hemorrhoids excised (Table 2). Moreover, the patients in the non-fluid-restriction group patients were significantly older and more likely to exhibit postoperative urinary retention.<sup>18</sup> This might be explained by the lower pain produced by the SH procedure and adequate pain relief after the surgery. In addition, postoperative pain can lead to urinary retention through inhibition of the micturition reflex via increased sympathetic nervous system activity.<sup>15</sup> We found here that fluid restriction itself did not cause pain in patients who underwent hemorrhoidopexy at

any time after the operation (Fig. 2), or increase the use of analgesic drugs (three patients in the non-fluid-restriction group and five in the fluid-restriction group;  $p = 0.702$ ; Table 2). Although overdistension of the bladder is a known risk factor for urinary retention, it did not appear to cause this in our study.<sup>19</sup>

Unexpectedly – as shown in Fig. 3 – there was no statistically significant difference in the average time to first urination between the two groups ( $700.04 \pm 455.03$  min in the non-fluid-restriction group and  $737.16 \pm 426.32$  min in the fluid-restriction group;  $p = 0.669$ ). This probably arose because all of our patients received intravenous anesthesia so the use of perioperative intravenous fluid was inevitable. Although the perioperative intravenous fluid volumes ( $218 \pm 87.66$  mL in EG and  $212 \pm 89.30$  mL in CG), with some oral intake, when combined with irritation from the anal wound, were enough to trigger urination. Therefore, fluid restriction had no effect on delaying urination in these patients.

Although spinal anesthesia with the patient in a prone jackknife position with detailed history taking by the anesthesiologist is thought to enable airway patency, all of our patients were subjected to intravenous anesthesia plus local anesthesia safely. Spinal anesthesia can dull bladder sensations and inhibit the voiding reflex.<sup>19</sup> This could also explain why our patients had lower urinary retention rates after the operation.

In terms of postoperative complications, there were two cases of delayed bleeding, and both of them were in the non-fluid-restriction group. According to previous studies, such bleeding occurs either immediately or 4-10 days after surgery.<sup>20,21</sup> Delayed bleeding in our two cases occurred at 10 and 14 days after surgery, but we consider that it had no link with fluid restriction. Nisar et al. published a meta-analysis on 15 prospective randomized trials, and reported that the hemorrhoidopexy: hemorrhoidectomy ratio of postoperative rectal bleeding was 2.3 to 1.<sup>22</sup> In addition, in both of our cases with this complication we used a PPH stapler, which has been reported to lead to more postoperative bleeding than the EEA device.<sup>23</sup>

### Limitations

Although this was a prospective, double-blinded, randomized controlled trial, the strength of this study was limited by the small sample size and because it was a single-center experience. The low numbers of recruited patients might lead to low power analysis in this study. Further multiple-center trials are needed to test the validity of our findings.

### Conclusions

Fluid restriction after SH with or without partial external hemorrhoidectomy did not reduce postoperative pain, postpone the time to first urination, or decrease the risk of postoperative urinary retention. Therefore, fluid restriction after SH is unnecessary.

### Consent to Publish

Not applicable.

### Availability of Data and Materials

All the data regarding the findings are available within the manuscript.

### Authors' Contributions

Chun-Wei Yu: data collection and wrote the manuscript. Nien-Ying Tsai: data collection. Shu-Wen Jao: recruited the patients and did the operations. Chao-Yang Chen: recruited the patients and did the operations. Chia-Cheng Wen: recruited the patients and did the operations. Chien-Chang Kao: analyzed the reports of uroflowmetry and residual urine. Kuan-Ling Lin: data analysis and randomize the patients. Po-Hsien Wu: data collection. Yi-Chiao Cheng: designed the study and wrote the manuscript.

### Acknowledgement

We would like to acknowledge the data collection and patient treatment provided by the members of the Division of Colon and Rectal Surgery, Department of Surgery, and Division of Urological Surgery, Department of Surgery.

### Funding

This study was supported by the study project of Tri-Service General Hospital (ATSGH-C107-201). The funder had no role in study design and data analyze.

### Conflict of Interest

All the authors declare that no conflict of interest.

### Ethical Approval

This study was reviewed and approved by the Tri-Service General Hospital Institutional Review Board for human subjects (No. 2-106-05-063). This study also had trial registration with clinicalTrials.gov (Identifier: NCT04459039). The first registration was July 7<sup>th</sup> 2020.

## Informed Consent

Informed consent was obtained from all individual participants included in the study.

## References

- Ratto C, Campenni P, Papeo F, Donisi L, Litta F, Parello A. Transanal hemorrhoidal dearterialization (THD) for hemorrhoidal disease: a single-center study on 1000 consecutive cases and a review of the literature. *Techniques in Coloproctology* 2017;21(12):953-62.
- Riss S, Weiser FA, Schwameis K, et al. The prevalence of hemorrhoids in adults. *International Journal of Colorectal Disease* 2012;27(2):215-20.
- Geller EJ. Prevention and management of postoperative urinary retention after urogynecologic surgery. *International Journal of Women's Health* 2014;6:829.
- Prasad ML, Abcarian H. Urinary retention following operations for benign anorectal diseases. *Diseases of the Colon & Rectum* 1978;21(7):490-2.
- Agrawal K, Majhi S, Garg R. Post-operative urinary retention: review of literature. *World Journal of Anesthesiology* 2019;8(1):1-12.
- Bailey HR, Ferguson JA. Prevention of urinary retention by fluid restriction following anorectal operations. *Diseases of the Colon & Rectum* 1976;19(3):250-2.
- Longo A. Treatment of hemorrhoidal disease by reduction of mucosa and hemorrhoidal prolapse with a circular stapler suturing device: a new procedure. *Proceeding of the 6<sup>th</sup> World Congress of Endoscopic Surgery* 1998;1998:777-84.
- Correa-Rovelo JM, Tellez O, Obregón L, Miranda-Gomez A, Moran S. Stapled rectal mucosectomy vs. closed hemorrhoidectomy. *Diseases of the Colon & Rectum* 2002;45(10):1367-76.
- Ravo B, Amato A, Bianco V, et al. Complications after stapled hemorrhoidectomy: can they be prevented? *Techniques in Coloproctology* 2002;6(2):83-8.
- Mehigan BJ, Monson JR, Hartley JE. Stapling procedure for haemorrhoids versus Milligan-Morgan haemorrhoidectomy: randomised controlled trial. *The Lancet* 2000;355(9206):782-5.
- Zaheer S, Reilly WT, Pemberton JH, Ilstrup D. Urinary retention after operations for benign anorectal diseases. *Diseases of the Colon & Rectum* 1998;41(6):696-704.
- Kunitake H, Poylin V. Complications following anorectal surgery. *Clinics in Colon and Rectal Surgery* 2016;29(1):14-21.
- Gottesman L, Milson JW, Mazier WP. The use of anxiolytic and parasympathomimetic agents in the treatment of postoperative urinary retention following anorectal surgery. *Diseases of the Colon & Rectum* 1989;32(10):867-70.
- Jackson J, Davies P, Leggett N, et al. Systematic review of interventions for the prevention and treatment of postoperative urinary retention. *BJS Open* 2019;3(1):11-23.
- Toyonaga T, Matsushima M, Sogawa N, et al. Postoperative urinary retention after surgery for benign anorectal disease: potential risk factors and strategy for prevention. *International Journal of Colorectal Disease* 2006;21(7):676-82.
- Iusuf T, Sarbu V, Cristache C, Popescu R, Botea F, Panait L. Urinary complications after anorectal surgery. *Chirurgia* 2000;95(6):531-4.
- Petros JG, Bradley TM. Factors influencing postoperative urinary retention in patients undergoing surgery for benign anorectal disease. *The American Journal of Surgery* 1990;159(4):374-6.
- Kozol RA, Mason K, McGee K. Post-herniorrhaphy urinary retention: a randomized prospective study. *Journal of Surgical Research* 1992;52(2):111-2.
- Tammela T. Postoperative urinary retention--why the patient cannot void. *Scandinavian Journal of Urology and Nephrology Supplementum* 1995;175:75-7.
- Pescatori M, Gagliardi G. Postoperative complications after procedure for prolapsed hemorrhoids (PPH) and stapled transanal rectal resection (STARR) procedures. *Techniques in Coloproctology* 2008;12(1):7-19.
- Angelone G, Giardiello C, Prota C. Stapled hemorrhoidectomy. Complications and 2-year follow-up. *Chirurgia Italiana* 2006;58(6):753-60.
- Nisar PJ, Acheson AG, Neal KR, Scholefield JH. Stapled hemorrhoidectomy compared with conventional hemorrhoidectomy: systematic review of randomized, controlled trials. *Diseases of the Colon & Rectum* 2004;47(11):1837-45.
- Giuratrabocchetta S, Pecorella G, Stazi A, Tegon G, De Fazio M, Altomare D. Safety and short-term effectiveness of EEA stapler vs PPH stapler in the treatment of degree III haemorrhoids: prospective randomized controlled trial. *Colorectal Disease* 2013;15(3):354-8.



原 著

## 痔瘡環切術後需要限水嗎？一項前驅、雙盲、隨機對照試驗

游鈞偉<sup>1</sup> 蔡念螢<sup>2</sup> 饒樹文<sup>1</sup> 陳昭仰<sup>1</sup> 溫家政<sup>1</sup>  
高建璋<sup>3</sup> 林冠伶<sup>1</sup> 吳柏憲<sup>1</sup> 鄭屹喬<sup>1</sup>

<sup>1</sup>三軍總醫院 外科部 大腸直腸外科

<sup>2</sup>三軍總醫院 一般醫學部

<sup>3</sup>三軍總醫院 外科部 泌尿外科

常規痔瘡切除術易導致術後尿滯留，因此術後常需要限制水分攝取以降低併發症發生。然而，痔瘡環切術術後是否需要限水仍未有定論。

為此我們設計了一個雙盲、隨機對照試驗，將 55 位三、四級痔瘡病人分為兩組，在接受痔瘡環切術後 12 小時內，一組只能攝取 250 ml，而另一組給予 1000 ml 點滴，比較術後尿滯留及疼痛的比例。

研究結果發現，術後無論是否有水份攝取限制，疼痛分數及術後尿滯留發生比率均無顯著差異。

**關鍵詞** 痔瘡環切術、術後尿滯留、術後疼痛、限水。